

## Effects of estrogen replacement therapy on the lipoprotein profile in postmenopausal women with ESRD

ELIZABETH S. GINSBURG, BRIAN WALSH, LOUISE GREENBERG, DEBORAH PRICE, GLENN M. CHERTOW, and WILLIAM F. OWEN, JR.

Department of Obstetrics and Gynecology; Division of Endocrinology and Metabolism; and Renal Division, Department of Medicine; Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts USA

### Effects of estrogen replacement therapy on the lipoprotein profile in postmenopausal women with ESRD.

**Background.** Patients with ESRD have excessive cardiovascular morbidity and mortality. In postmenopausal women with normal renal function, estrogen replacement therapy decreases cardiovascular mortality by 50%, in part because of their beneficial effects on the lipoprotein profile. Because of similarities in the lipoprotein profile between healthy, postmenopausal women, and women with ESRD, we examined the effects of estrogen replacement on lipoproteins in 11 postmenopausal women with ESRD.

**Methods.** In a randomized, placebo-controlled crossover study (8 week treatment arms) using 2 mg daily of oral, micronized estradiol, 11 postmenopausal women with ESRD were treated. Neither baseline lipid nor lipoprotein abnormalities were used as entry criteria for study participation.

**Results.** Blood estradiol levels were  $19 \pm 4$  with placebo and  $194 \pm 67$  pg/ml ( $P = 0.024$ ) with estradiol treatment. Total HDL cholesterol concentrations increased from  $52 \pm 19$  mg/dl to  $61 \pm 20$  mg/dl (16%), with placebo and estradiol treatments, respectively ( $P = 0.002$ ). Apolipoprotein A<sub>1</sub> increased by 24.6% ( $P = 0.0002$ ) with estradiol intervention. HDL<sub>2</sub> concentrations were  $19 \pm 13$  with placebo and  $24 \pm 16$  with estradiol treatment ( $P = 0.046$ ). There were no differences in total or LDL cholesterol, other lipoprotein fractions including Lp(a), and triglycerides with 2 mg daily estradiol treatment. No significant side effects were observed.

**Conclusions.** Therefore, using standard dosage regimens for estrogen replacement therapy in postmenopausal women with ESRD, HDL cholesterol is increased to an extent that would be expected to improve their cardiovascular risk profile. Further studies are needed to assess whether estrogen replacement therapy decreases the incidence or severity of cardiovascular disease in ESRD patients to a similar degree compared with other women.

In contrast to age- and sex-matched controls with normal renal function, cardiovascular morbidity and mortality are

excessive in patients with end-stage renal disease (ESRD) [1–3]. Arguably, patients with ESRD undergo accelerated atherosclerosis [4–6], which is a consequence of deleterious qualitative and quantitative lipid abnormalities that accompany renal insufficiency [7–9]. Common lipid abnormalities that are of potential clinical relevance for patients with renal failure are: (1) diminished high density lipoprotein (HDL) cholesterol and apolipoprotein A<sub>1</sub> concentrations; (2) normal or elevated low density lipoprotein (LDL) concentrations; (3) hypertriglyceridemia with increased very low density lipoproteins (VLDL); and (4) increased lipoprotein (a) [Lp(a)] concentrations [6–13]. However, with the notable exception of Lp(a), which is an independent risk factor for atherosclerotic vascular disease in patients with normal renal function [14–18] and ESRD [12], most investigators have had difficulty establishing a correlation between the lipoprotein profile and cardiovascular events in ESRD patients [1, 19, 20].

Reproductive hormones such as estrogen have a profound influence on the lipoprotein profile of both men and women. Supportive evidence for the role of sex hormones in lipid metabolism is derived from the observation of gender differences in LDL and HDL levels with age [21, 22], the effect of the menstrual cycle on plasma LDL and HDL concentrations [23, 24], and the effect of oral estrogens or progestins on the lipoprotein profile of men with prostate cancer and postmenopausal women [25–29]. For example, postmenopausal women have higher total and LDL cholesterol concentrations, and lower HDL cholesterol levels, than premenopausal women. Postmenopausal women also have an increased rate of cardiovascular events. The deleterious lipid profile and cardiovascular outcome has been ascribed to the menopausal loss of estrogen [20, 21, 30, 31]. In contrast, postmenopausal women receiving estrogen replacement therapy have half the relative risk of death due to cardiovascular disease as do postmenopausal women who do not use estrogen. Some investigators have advocated that the cardiovascular risk

**Key words:** cardiovascular disease, lipoproteins, cholesterol, estrogen, lipids, end-stage renal disease, menopause.

Received for publication January 13, 1998

and in revised form April 22, 1998

Accepted for publication April 23, 1998

© 1998 by the International Society of Nephrology

reduction from estrogen replacement therapy is a consequence of their effect on the deleterious lipoprotein profile of being postmenopausal [32–34]. Specifically, estrogen decreases LDL cholesterol by augmenting hepatic LDL catabolism. HDL cholesterol is increased, especially HDL<sub>2</sub>. Other quantitative and qualitative effects of estrogen include: increases in apolipoprotein A<sub>1</sub> in HDL, but not A<sub>2</sub>; increased triglyceride in VLDL; and reduced Lp(a), both total and isoform specific [20, 25, 27].

It is unclear how vigorously to treat hyperlipidemia in ESRD [6–8]. The frequency and severity of complications associated with the long term use of some conventional lipid lowering agents in patients with ESRD may render their preemptive, routine use unwise [8]. Based on the effects of estrogen replacement therapy on the concentrations and metabolism of plasma lipoproteins in healthy postmenopausal women, the correction in the profile of lipoproteins is remarkably selective for the pathobiologic pattern of lipid abnormalities seen in patients with ESRD. Therefore, the use of estrogen replacement may be a novel preemptive pharmacological intervention for the lipid abnormalities in women with ESRD. We structured this study to address the hypothesis that estrogen replacement therapy will favorably improve the lipoprotein profile in postmenopausal women with ESRD undergoing maintenance hemodialysis.

## METHODS

This study was approved by the Human Research Committee of the Brigham and Women's Hospital. All subjects provided informed consent. Eligible subjects were women  $\geq 45$  years old who had undergone hemodialysis for  $\geq$  six months, and no menstrual periods for  $\geq$  one year. Inclusion criteria were stable hematocrit  $\geq 22\%$ , normal hepatic transaminases, and a follicle stimulating hormone greater than 40 IU/ml. Exclusion criteria were a history of a cerebrovascular accident, breast cancer, or current diabetes mellitus requiring insulin therapy. Neither baseline lipid nor lipoprotein abnormalities were used as entry criteria for study participation. No patient was on estrogen replacement therapy for six weeks prior to entry into the study.

### Study protocol

All subjects underwent an initial history and physical examination, including a screening mammogram. Seven subjects entered a lead-in study evaluating the estradiol levels attained on 1 mg oral estradiol daily for four weeks. Because of concerns about the achievement of excessive sex hormone levels, and a desire to minimize patient risks secondary to excessive estrogen exposure, the lead-in study was performed to assess the impact of a lower dose of estrogen replacement therapy. After completing the four week lead-in study, the subjects took medroxyprogesterone acetate, 10 mg daily for one week and then underwent a four week wash-out period. Medroxyprogesterone acetate

was included in this and the subsequent protocol to counteract the uterine effects of unopposed estrogen stimulation. The dose and duration of medroxyprogesterone acetate therapy were conventional [25]. Based on these results, thirteen subjects entered a double-blinded cross-over study using oral micronized estradiol (2 mg) and a placebo packaged in identical blisterpacks (supplied by Mead-Johnson, Bristol Myers-Squibb). The women were initially randomized to receive estrogen daily (or placebo) for eight weeks, and after a wash-out period of four weeks, they received the other therapy for eight weeks. Prior to receiving the first intervention, subjects who had not participated in the lead-in study took medroxyprogesterone acetate, 10 mg daily for one week, and then began the first treatment arm four weeks later. After eight weeks of treatment with estrogen or placebo, all subjects took medroxyprogesterone acetate, 10 mg daily for one week. Fasting blood samples were obtained: (1) two days during the week prior to the first intervention (estrogen or placebo), baseline samples, (2) two days during the fourth week of the intervention, and (3) three days during the last (eighth) week of the intervention. After completion of the second treatment arm, women with intact uteri took medroxyprogesterone acetate, 10 mg daily for one week.

### Blood drawing, sample processing, and assays

All blood samples were drawn from the hemodialysis angioaccess at the time of routine predialysis blood drawing with the subject fasting overnight. Samples were refrigerated, centrifuged at 2,500 rpm, aliquotted, and stored at  $-70^{\circ}\text{C}$ . Estradiol was assayed using the immunofluorescent Delphia system (Wallac, Gaithersburg, MD, USA). The intraassay coefficient of variation for estradiol concentration is 10% from 0 to 99 pg/ml, 4.5% from 100 to 299 pg/ml, and 2.7% at  $\geq 300$  pg/ml. The interassay coefficient of variation is 4.6% from 0 to 300 pg/ml, 2.7% from 301–1000 pg/ml and 5.3% at  $>1000$  pg/ml. There is  $<1\%$  cross-reactivity with estrone. Estrone was assayed by radioimmunoassay using a commercial kit from Wein (Succasunna, NJ, USA). The lower limit of sensitivity is 20 pg/ml. At mean estrone levels of 80, 202 and 466 pg/ml the mean intraassay coefficients of variation are 8.4%, 5.6%, and 6.2%, respectively. At mean estrone levels of 53, 205 and 497 pg/ml, the mean coefficients of variation were 13.5%, 5.6% and 2.6%, respectively. Sex hormone binding globulin (SHBG) and serum albumin concentrations (BCG method) were measured by conventional means. Estrone, albumin and SHBG concentrations were measured on samples pooled from the three samples drawn the last week of each eight week treatment arm.

Blood samples from each subject were analyzed in one batch at the end of the study in the Lipid Research Laboratory of the Nutrition Department, Harvard School of Public Health (Boston, MA, USA). This study was standardized for cholesterol and HDL measurements as

specified by the Centers for Disease Control and the National Heart, Lung and Blood Institute. Samples from each blood draw were measured separately, and the results from each week (that is, results of three blood draws during week 8) were averaged. VLDL was prepared from plasma by overlaying 0.5 ml of 0.9% sodium chloride over 0.5 ml of plasma, and spinning in a Beckman Type 25 rotor (Beckman Instruments, Palo Alto CA, USA) at 25,000 rpm for 16 hours. The VLDL fraction was separated from the LDL and HDL fractions by tube slicing. HDL and HDL<sub>3</sub> were sequentially separated by precipitation with dextran sulfate and magnesium chloride [35]. HDL<sub>2</sub> was obtained by subtraction of HDL<sub>3</sub> from the total HDL. Cholesterol and triglycerides were measured with enzymatic reagents and quantified photometrically using a COBAS Mira Plus auto-analyzer (Roche Diagnostics Systems, Belleville, NJ, USA). Cholesterol, VLDL, the combined LDL and HDL fractions, HDL and HDL<sub>3</sub> were measured in whole plasma. Triglycerides and apolipoprotein B were measured in whole plasma and in the VLDL fraction. Apolipoprotein B, apolipoprotein A<sub>1</sub> and Lp(a) were measured by immunoturbidimetry with rabbit antiserum, [36] obtained from Incstar (Stillwater, MN, USA) using the COBAS Mira Plus auto-analyzer. The coefficients of variation for blinded control specimens were as follows: 2.4% for cholesterol, 3.3% for HDL cholesterol, 2.8% for HDL<sub>3</sub> cholesterol, 2.2% for apolipoprotein A<sub>1</sub>, 2.0% for apolipoprotein B, and 3.0% for Lp(a).

### Statistical analysis

Duplicate measurements were averaged and weighted as a single test. Pre- and post-estrogen exposure lipoprotein and hormone parameters were compared using the Wilcoxon rank sum test. Comparisons were considered statistically significant at  $P < 0.05$ . Non-parametric tests, like the Wilcoxon rank sum, tend to be far more conservative with regard to type I error than corresponding parametric tests (such as, Student's *t*-test). Analyses were conducted using SAS (The SAS Institute, Cary, NC, USA).

### RESULTS

The principal rationale for the lead-in study of four weeks with 1 mg micronized oral estradiol daily was to assess its impact on estrogen levels and to minimize patient risks secondary to excessive estrogen exposure, not as an intervention to effect the lipoprotein profile. Seven subjects received 1 mg oral estradiol daily. One subject withdrew because of ill health during fasting prior to blood sampling at the time of hemodialysis. The remaining six subjects completed the lead-in study, and three of these women participated and completed the blinded, crossover portion of the study. Thirteen subjects total were randomized to the blinded crossover trial, but serum samples from one subject were not able to be analyzed because of a handling mishap. Twelve subjects completed the study. One subject withdrew

**Table 1.** Demographic profile of subjects completing randomized trial

Subject	Age years	Etiology of ESRD	Length of renal replacement Rx years	Estimated dry weight kg
1	53	hypertension	4	59
2	62	polycystic kidney disease	2	54.5
3	56	hypertension	23	63
4	72	hypertension	0.7	55.5
5	45	focal glomerulosclerosis	0.5	70.5
6	59	hereditary nephritis	9.5	70
7	65	diabetes mellitus	0.5	66.4
8	75	diabetes mellitus	1.3	82
9	63	unknown	3.0	56.5
10	70	hypertension	10	57.5
11	58	hereditary nephritis	0.5	66.5

**Table 2.** Hormonal and lipid parameters with four weeks of estradiol

	Baseline value	Value at week 4	P value
Estradiol pg/ml	21 ± 6	91 ± 19	0.002
Total cholesterol mg/dl	222.7 ± 29.9	247.0 ± 32.7	0.001
Total HDL cholesterol mg/dl	46.5 ± 4.9	55.6 ± 5.5	0.029
LDL cholesterol mg/dl	45.5 ± 7.3	43.4 ± 7.2	0.549
HDL <sub>2</sub> cholesterol mg/dl	8.9 ± 2.8	12.8 ± 4.1	0.052
HDL <sub>3</sub> cholesterol mg/dl	37.6 ± 3.7	42.8 ± 2.8	0.087
Triglycerides mg/dl	227.4 ± 36.3	217.2 ± 36.3	0.549

Abbreviations are: LDL, low density lipoprotein; HDL, high density lipoprotein.

Results are from six women receiving 1 mg per day of estradiol

because she underwent cadaveric renal transplantation during the trial. Another subject experienced vaginal bleeding due to a benign vaginal lesion that was uneventfully excised; she completed the study. Demographic information on the 11 subjects who completed the trial is provided in Table 1. Their mean age was  $61.4 \pm 2.7$  years ( $\pm$  SEM); eight of the women were African American, one was Hispanic, and two were white. Two women had non-insulin dependent diabetes mellitus.

### Lipoprotein profiles

Lipoprotein and estradiol measurements were obtained from the six women who completed the preliminary intervention trial during which they ingested 1 mg estradiol daily for four weeks. Table 2 provides the change in the lipid profile for these subjects. During the period of estradiol ingestion, estradiol levels increased approximately 4.5-fold. In parallel, total cholesterol, total HDL cholesterol, and the HDL<sub>2</sub> cholesterol fraction increased significantly. No significant change was observed in triglycerides.

Hormonal parameters from the placebo controlled, cross-over intervention trial are provided in Table 3. Higher estradiol levels were achieved with 2 mg per day of estradiol, in comparison to 1 mg daily. In comparison to eight weeks of placebo therapy, statistically significant increases in estradiol, its metabolite estrone, and the

**Table 3.** Hormonal levels with eight weeks of therapy

Hormonal parameter	Intervention			Reference range
	Placebo	Estradiol <sup>a</sup>	P value	
Estradiol pg/ml	19 ± 4	194 ± 67	0.024	<30
Estrone pg/ml	24 ± 6	684 ± 186	0.006	15–55
Albumin mg/dl	4033 ± 118	3947 ± 84	0.220	3900–5400
Sex hormone binding globulin nmol/liter	61 ± 7	94 ± 9	0.0006	8–85

<sup>a</sup> Results are from eleven women receiving 2 mg per day of estradiol


**Table 4.** Serum lipoprotein levels with eight weeks of therapy (mean ± SEM)

Lipoprotein mg/dl	Intervention			Reference range
	Placebo	Estradiol <sup>a</sup>	P value	
<b>VLDL</b>				
Cholesterol	22.6 ± 19.6	24.2 ± 20.8	0.643	<20
Triglycerides	78.7 ± 62.4	92.4 ± 81.2	0.319	NE <sup>b</sup>
Apolipoprotein B	17.0 ± 13.4	17.9 ± 13.5	0.647	NE
<b>LDL</b>				
Cholesterol	131.0 ± 56.2	129.1 ± 49.1	0.837	<130
Apolipoprotein B	55.9 ± 21.7	57.1 ± 17.7	0.761	NE
<b>HDL</b>				
Cholesterol	52.2 ± 19.4	60.6 ± 19.6	0.002	>35
<b>HDL<sub>2</sub></b>				
Cholesterol	18.5 ± 12.7	23.8 ± 15.9	0.045	NE
<b>HDL<sub>3</sub></b>				
Cholesterol	33.7 ± 9.7	36.8 ± 8	0.106	NE
Apolipoprotein A <sub>1</sub>	149.7 ± 42.2	186.6 ± 39.1	0.0002	115–220
Total cholesterol	205.4 ± 71.8	211.4 ± 56.6	0.538	<200
Total triglycerides	164.1 ± 96.2	190.1 ± 103.4	0.176	<130
Lp(a)	36.8 ± 33.3	36.8 ± 29.2	0.987	NE

<sup>a</sup> Each of eleven women received 2 mg per day of estradiol

<sup>b</sup> NE, normal range has not been established

steroid-inducible SHBG were observed. Total HDL cholesterol concentration increased in all subjects (Table 4 and Fig. 1A). The increase of the mean values was 16.1%. Total HDL cholesterol concentration increased from 52.2 ± 19.4 mg/dl after placebo treatment to 60.6 ± 19.6 mg/dl (mean ± SEM;  $P = 0.002$ ) following estradiol. The fraction of HDL that accounted for the majority of the increase in total HDL cholesterol was HDL<sub>2</sub>, which increased by 28.6% ( $P = 0.046$ ), whereas HDL<sub>3</sub> cholesterol increased by 9.2%, a difference that did not reach statistical significance (Table 4). With eight weeks of estradiol treatment, nine of eleven women had increases in HDL<sub>2</sub> cholesterol, and eight women had increases in HDL<sub>3</sub> cholesterol. LDL cholesterol was unchanged for all subjects. There was no significant association between the increase in HDL cholesterol concentration and the concentrations of estradiol or estrone. Apolipoprotein A<sub>1</sub> increased by 24.6%, from 149.7 ± 42.2 mg/dl to 186.6 ± 39.1 mg/dl ( $P = 0.0002$ ). Lp(a) was not significantly affected. Although the total triglyceride concentration increased by 15.8% with daily estradiol ingestion, the elevation compared to placebo therapy was not significant.



**Fig. 1.** Individual total HDL cholesterol concentration (mg/dl) in 11 women with completing eight weeks of placebo and with eight weeks of estradiol (2 mg per day).

## DISCUSSION

Low HDL cholesterol concentrations are often, but not uniformly observed in patients with chronic renal insufficiency, especially the HDL<sub>2</sub> subfraction [37]. In the present study, HDL cholesterol concentrations were within the normal range. However, oral micronized estradiol increased the serum total HDL cholesterol concentration by an average of 8.4 mg/dl (16.1%), and increased apolipoprotein A<sub>1</sub> by 24.6% compared with placebo treatment. Previous studies in patients with normal renal function suggest that comparable increases in total HDL cholesterol can be expected to decrease the risk of cardiovascular disease by approximately 50% [38]. Even the 1 mg daily dose of estradiol that was administered during the lead-in portion of the current intervention trial increased HDL by 20% after only four weeks. This is a level of response that would also be expected to significantly decrease the incidence of coronary artery disease. The triglyceride concentration was not significantly increased by low dose estrogen replacement therapy, but the lack of statistical significance may be due in part to the small sample size in the setting of large inter-individual variations in triglyceride concentrations. The mean triglyceride levels in this subject group were not out of the normal range, however. Hypertriglyceridemia



seen in hemodialysis patients is thought to be due to defective triglyceride removal secondary to inhibition of lipoprotein lipase activity [39]. The doses of estrogen used in the intervention reported herein had no significant effect on plasma Lp(a) or LDL cholesterol levels. Insufficient power (that is, large type 2 error) is an important consideration in interpreting any apparent lack of effect, since size (and scatter) may limit the ability to detect differences among groups.

Previous investigations have demonstrated that basal estradiol concentrations as well as those attained after oral estradiol ingestion are greater in postmenopausal women with ESRD than in body-mass index matched controls, probably due to decreased clearance of endogenous and exogenous estradiol [40]. Although no adverse effects were observed in the present study, a larger number of subjects with a longer period of estrogen ingestion is needed to validate this observation. Because of the absence of correlation between the increase in estradiol and estrone concentrations and the increase in HDL cholesterol, it is unlikely that extraordinarily high sex hormone concentrations are needed to improve the HDL:LDL ratio. In that the biologic effect of 2 mg daily of oral estradiol is not much different than that observed with 1 mg daily, but blood estradiol levels were lower with 1 mg daily oral estradiol (194 vs. 91 pg/ml for 2 vs. 1 mg/day, respectively), we advocate the use of the lower estradiol dose for future clinical intervention trials. This will minimize the risks associated with higher hormone levels.

Not all ESRD patients have abnormal lipoprotein profiles. Moreover, the incidence of death due to cardiovascular disease in the ESRD population is very high regardless of the lipoprotein profiles found. Therefore, other factors may contribute to this excessive cardiovascular mortality, and/or the lipoprotein profile is an incomplete surrogate of the deleterious effect of hyperlipidemia in ESRD. Similarly, in healthy postmenopausal women, estrogen replacement therapy is associated with a decrease in the risk of death due to cardiovascular disease by 50% [41]. However, this effect appears to be partly independent of the individuals' lipoprotein profile, and is much greater than would be predicted by the change in the lipoprotein profile alone. Furthermore, based on longitudinal studies of healthy women with angiographically demonstrable coronary artery stenosis, simply having received estrogen replacement therapy at any time reduces the risk of death by 10 to 27% in comparison to nonusers [42]. Therefore, although the change in the lipoprotein profile was moderate in this study as in studies of women with normal renal function, a similar experience in postmenopausal women with ESRD may result in substantial benefit with reductions in deaths due to cardiovascular disease.

Many treatments for hyperlipidemia have been evaluated in ESRD including gemfibrozil [43], probucol [44], pan-tethine [45], and LDL-pheresis [46]. Lovastatin and simva-

statin treatment in ESRD patients on hemodialysis increases HDL cholesterol and apolipoprotein A<sub>1</sub> and decreases triglycerides [47]. However, whether these changes reduce the incidence of heart disease has not been established. Furthermore, consideration must be made of the major side effects associated with the use of many of the conventional lipid lowering agents. For example, HMG CoA reductase inhibitors have major routes of renal excretion. When used alone, but especially in combination therapy with other lipid lowering agents, there is an increased incidence of rhabdomyolysis and elevated hepatic transaminases. Similar side effects have been observed with gemfibrozil, as well as enhanced sensitivity to oral anticoagulants. Probucol is relatively unsafe in patients with conduction system abnormalities. Some data suggests that modification of dietary fat can significantly decrease triglycerides in hemodialysis patients [48]. However, if elevations of HDL cholesterol seen with gemfibrozil decreases cardiovascular risk [43], similar benefit may be achieved from the use of estrogen replacement therapy with fewer side effects. Added cardiovascular benefits from estrogens include their ability to dilate coronary arteries [49], decrease coronary resistance [50], and increase coronary perfusion.

The data presented herein indicate that the use of estrogen replacement therapy results in increases of HDL cholesterol and apolipoprotein A<sub>1</sub> that are comparable to those achieved in healthy postmenopausal women. In the present study, baseline lipid or lipoprotein abnormalities were not used as entry criteria. A principal reason for this approach is that in this early intervention trial, we were uncertain what benchmark abnormalities to use. Instead, we elected to examine the effect of estrogen replacement therapy on a random sample of postmenopausal women with ESRD, as was done in a previous landmark trial in healthy postmenopausal women [25]. Furthermore, the strategy of including ESRD patients, indifferent of the severity of their lipid abnormalities, permitted us to profile the type and severity of baseline lipid and lipoprotein derangements in a random sample of postmenopausal women with ESRD. The findings reported herein may not be reproducible in some postmenopausal women with an alternative lipid or lipoprotein profile. It is unknown whether patients with greater derangement in lipoprotein parameters would achieve the same, greater, or lesser effects with estrogen replacement as the subjects in the present study. Furthermore, this study has not addressed the ultimate question of whether improvements in lipoprotein parameters will lead to comparable reductions in cardiac mortality. Larger clinical trials are needed to answer these questions, and to determine whether estrogen replacement therapy should be conventional care for postmenopausal women with ESRD. In the design of future trials with estrogen replacement therapy, consideration

should be made of their potential to improve other comorbid conditions in postmenopausal women such as prevention of estrogen-deficiency related osteoporosis [51, 52], symptoms of urogenital atrophy [53], improved skin elasticity [54] and decreased wrinkling [55], and a decreased risk of Alzheimer's disease [56–59]. In patients with ESRD, platelet dysfunction is improved [60, 61]. These potential benefits must be balanced against the possible increased risks of breast cancer [62, 63], non-fatal venous thromboembolism [64], and cholelithiasis [65]. The increase risk of thromboembolic disease may be a consequence of the increase in clotting factors observed with oral, but not transdermal, estrogen replacement therapy [66, 67].

## ACKNOWLEDGMENTS

This study was supported by grants from Baxter, Inc. and Mead-Johnson, Inc. Partial funding for this work (to W.F.O.) was by grant #DK520521 of the National Institute of Diabetes, Digestive, and Kidney Disease.

Reprint requests to William F. Owen, Jr., M.D., Brigham and Women's Hospital, Dialysis Unit Administrative Office, 75 Francis Street, Boston, Massachusetts 02115, USA.  
E-mail: wfowen@bics.bwh.harvard.edu

## REFERENCES

- ROSTAND SG, KIRK KA, RUTSKY EA: Relationship of coronary risk factors to hemodialysis-associated ischemic heart disease. *Kidney Int* 22:304–308, 1982
- NATIONAL CENTER FOR HEALTH STATISTICS: *Health, United States, 1990*. Hyattsville, Public Health Service, 1991
- US RENAL DATA SYSTEM: *USRDS 1990 Annual Data Report*. Bethesda, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 1990
- LINDER A, CJARRA B, SHERRARD D, SCRIBNER BH: Accelerated atherosclerosis in prolonged maintenance hemodialysis. *N Engl J Med* 290:697–701, 1974
- GREEN D, STONE NJ, KRUMLOVSKY A: Putative atherogenic factors in patients with chronic renal failure. *Prog Cardiovasc Dis* 62:133–144, 1983
- FOLEY RN, PARFREY PS, HARNETT JD, KENT GM, MARTIN CJ, MURRAY DC, BARRE PE: Clinical and echocardiographic disease in patients starting end-stage renal disease therapy. *Kidney Int* 47:186–192, 1995
- ATTMAN P-O, ALAUPOVIC P: Lipid abnormalities in chronic renal insufficiency. *Kidney Int* 39(Suppl):S16–S32, 1991
- APPEL G: Lipid abnormalities in renal disease. *Kidney Int* 39:169–183, 1991
- WANNER C, FROMMERZ K, HORL WH: Hyperlipoproteinemia in chronic renal failure: Pathophysiological and therapeutic aspects. *Cardiology* 78:202–217, 1991
- PARRA HJ, MEXDOUR H, CACHERA C, DRACON M, TACOUTE A, FRUCHART JC: Lp(a) lipoprotein in patients with chronic renal failure treated by hemodialysis. (abstract) *Clin Chem* 33:721, 1987
- HAFFNER SM, GRUBRE KK, ALDRETE G, MORALES PA, STERN MP, TUTTLE KR: Increased lipoprotein (a) concentrations in chronic renal failure. *J Am Soc Nephrol* 3:1156–1162, 1992
- CRESSMAN MD, HEYKA RJ, PAGANINI EP, O'NEAL J, SKIBINSKI CI, HOFF HF: Lipoprotein (a) is an independent risk factor for cardiovascular disease in hemodialysis patients. *Circulation* 86:475–482, 1992
- KAYSEN GA: Hyperlipidemia of chronic renal failure. *Blood Purif* 12:60–67, 1994
- KOSTNER GM, AVOGARO P, CAZZOLATO E, MARTH E, BITTOLO-BON G, QUINCI GB: Lipoprotein Lp (a) and the risk for myocardial infarction. *Atherosclerosis* 38:51–61, 1986
- RHOADS GG, DAHLEN G, BERG K, MORTON NE, DANNENBERG AL: Lp(a) lipoprotein as risk factor for myocardial infarction. *J Am Med Assoc* 256:2540–2544, 1986
- HOFF HF, BECK GJ, SKIBINSKI CI, JURGENS G, O'NEIL J, KRAMRE J, LYTLE B: Serum Lp (a) level as a predictor of vein graft stenosis after coronary bypass surgery in patients. *Circulation* 77:2138–2144, 1988
- SCANU AM, FLESS GM: Lipoprotein (a): Heterogeneity and biological relevance. *J Clin Invest* 85:1709–1715, 1990
- SAND KAMP M, FUNKE H, SCHULTE H, KOHLER E, ASSMAN G: Lipoprotein (a) is an independent risk factor for myocardial infarction at a young age. *Clin Chem* 36:20–23, 1990
- PONTICELLI C, BARBI G, CATALUPPI A, DONATI C, ANNONI G, BRANCACCI D: Lipid abnormalities in maintenance dialysis patients and renal transplant recipients. *Kidney Int* 13:572–578, 1978
- VINCENTI F, AMEND WJ, ABELE J, FEDUSKA NJ, SALVATIERRA O: The role of hypertension in hemodialysis-associated atherosclerosis. *Am J Med* 68:363–369, 1980
- The Lipid Research Clinics' Population Studies Data Book. The Prevalence Study*. Bethesda, US Department of Health and Human Services, Public Health Service, National Institutes of Health, NIH Publication No. 80–1572, 1980
- SACHS FM, WALSH BW: The effects of reproductive hormones on serum lipoproteins: Unresolved issues in biology and clinical practice. *Ann NY Acad Sci* 592:272–285, 1990
- KIM HJ, KALKHOFF RK: Changes in lipoprotein composition during the menstrual cycle. *Metabolism* 28:663–668, 1979
- MATTSON LA, SILFVERSTOLPE G, SAMSOIE G: Lipid composition of serum lipoproteins in relation to gonadal hormones during the normal menstrual cycle. *Eur J Obstet Gynecol Reprod Biol* 17:327–335, 1984
- WALSH BW, SCHIFF I, ROSNER B, GREENBERG L, RAVNIKAR V, SACKS FM: Effects of postmenopausal estrogen replacement of the concentrations and metabolism of plasma lipoproteins. *N Engl J Med* 325:1196–1204, 1991
- FARISH E, HART DM, GRAY CE, BEASTALL G, FLETCHER CD, LINSAY R: Effects of treatment with oestradiol/levonorgestrel on bone, lipoproteins and hormone status in postmenopausal women. *Clin Endocrinol* 31:607–615, 1989
- SOMA M, MESCHIA M, GHANEM K: Plasma Lp (a) concentration after oestrogen and progestagen in postmenopausal women. (letter) *Lancet* 337:612, 1991
- HENRIKSSON P, ANGELIN B, BERGLUND L: Hormonal regulation of serum Lp (a) levels. Opposite effects after estrogen treatment and orchidectomy in males with prostate carcinoma. *J Clin Invest* 89:1166–1171, 1992
- WALSH BW, SACHS FM: The effects of low dose oral contraceptives on very-low density and low density lipoprotein metabolism. *J Clin Invest* 91:2126–2132, 1993
- KANNEL WB, HJORTLAND M, MCNAMARA PM, GORDON T: Menopause and risk of cardiovascular disease. The Framingham Study. *Ann Int Med* 85:447–452, 1976
- GORDON T, KANNEL WB, HJORTLAND MD, MCNAMARA PM: Menopause and risk of cardiovascular disease. The Framingham Study. *Ann Int Med* 89:157–161, 1978
- HENDERSON BE, RSS RK, PAGANINI-HILL A, MACK TM: Estrogen use and cardiovascular disease. *Am J Obstet Gynecol* 154:1181–1186, 1986
- BUSH TL, BARRETT-CONNOR E, COWAN LD: Cardiovascular mortality and noncontraceptive use of estrogen in women: Results from the Lipid Research Clinic Program Follow-up Study. *Circulation* 75:1102–1109, 1987
- STAMPER MJ, COLDITZ GA, WILLETT WC: Postmenopausal estrogen therapy and cardiovascular disease: Ten year follow-up from the Nurses' Health Study. *N Engl J Med* 325:756–762, 1991
- BACKORIK PS, ALBERS JJ: Precipitation methods for quantification of lipoproteins, in *Plasma Lipoproteins* (part B, vol 129), *Methods in Enzymology*, edited by ALBERS JJ, SEGREST JP, Orlando, Academic Press, 1986, pp 78–100
- FINLEY PR, WILLIAMS JR, BYERS JM: Immunochemical determination of human immunoglobulins with a centrifugal analyzer. *Clin Chem* 22:1037–1041, 1976
- WANNER C, FROMMERZ K, HORL WH: Hyperlipoproteinemia in chronic renal failure. Pathophysiological and therapeutic aspects. *Cardiology* 78:2020–2027, 1991
- MANNINEN V, ELO MO, FRICK MH: Lipid alterations and decline in

- the incidence of coronary heart disease in the Helsinki Heart Study. *J Am Med Assoc* 260:641–651, 1988
39. CATTRAN DC, FENTON SSA, WILSON DR, STEINER G: Defective triglyceride removal in lipemia associated with peritoneal dialysis and haemodialysis. *Ann Int Med* 85:29–33, 1976
  40. GINSBURG E, OWEN WF, GREENBERG LM, SHEA BF, LAZARUS JM, WALSH B: Estrogen absorption and metabolism in postmenopausal women with end stage renal disease. *J Clin Endo Metab* 81:4414–4417, 1996
  41. STAMPFER MJ, COLDITZ GA, WILLETT WC: Postmenopausal estrogen therapy and cardiovascular disease. Ten year follow-up from the Nurses' Health Study. *N Engl J Med* 325:756–762, 1991
  42. SULLIVAN JM, VANDER ZWAAG R, HUGHES JP, MADDOCK V, KROETZ FW, RAMANATHAN KB, MIRIVIS DM: Estrogen replacement and coronary artery disease. Effect on survival in postmenopausal women. *Arch Intern Med* 150:2557–2562, 1990
  43. ELISAF MS, DARDAMANIS MA, PAPAGALANIS ND, SIAMOPOULOS KC: Lipid abnormalities in chronic uremic patients. Response to treatment with gemfibrozil. *Scand J Urol Nephrol* 27:101–108, 1993
  44. FIORINI F, PATRONE E, CASTELLUCCIO A: Clinical investigation on the hypolipidemic effect of simvastatin versus probucol in hemodialysis patients. *Clin Ther* 145:213–217, 1994
  45. CORONEL F, TORNERO F, TORRENTE J, NARANJO P, DE OLEO P, MACIA M, BARRIENTOS A: Treatment of hyperlipidemia in diabetic patients on dialysis with a physiological substance. *Am J Nephrol* 11:32–36, 1991
  46. BOSCH T, THIERY J, GURLAND HJ, SEIDEL D: Long-term efficiency, biocompatibility, and clinical safety of combined simultaneous LDL-apheresis and haemodialysis in patients with hypercholesterolaemia and end-stage renal failure. *Nephrol Dial Transplant* 8:1350–1358, 1993
  47. WANNER C, HORL WH, LULEY CH, WIELAND H: Effects of HMG-CoA reductase inhibitors in hypercholesterolemic patients on hemodialysis. *Kidney Int* 39:754–760, 1991
  48. WAHLQVIST ML, HURLEY BP: Hyperlipoproteinemia and dietary fat modification in haemodialysis and renal transplant patients. *Med J Aust* 2:207–208, 1977
  49. COLLINS P, ROSANO GM, SARREL PM, ULRICH L, ADAMOPOULOS S, BEALE CM, MCNEILL JG, POOLE-WILSON PA: 17-beta estradiol attenuates acetylcholine-induced coronary arterial constriction in women but not men with coronary heart disease. *Circulation* 92:24–30, 1995
  50. GILLIGAN DM, AUYYUMI AA, CANNON RO 3<sup>rd</sup>: Effects of physiological levels of estrogen on coronary vasomotor function in postmenopausal women. *Circulation* 89:2545–2551, 1994
  51. FELSON DT, SHANG Y, HANNAN MT: The effect of postmenopausal estrogen therapy on bone density in elderly women. *N Engl J Med* 329:1141–1146, 1993
  52. CAULEY JA, SEELEY DG, ENSRUD K: Estrogen replacement therapy and fractures in older women. *Ann Intern Med* 122:9–16, 1995
  53. CARR BR, MACDONALD PC: Estrogen treatment of postmenopausal women. *Adv Intern Med* 28:491, 1983
  54. HENRY F, PIERARD-FRANCHIMONT C, CAUWENBERGH G, PIERARD GE: Age related changes in facial skin contours and rheology. *J Am Geriatr Soc* 45:220–222, 1997
  55. DUNN LB, DAMESYN M, MOORE AA, REUBEN DB, GREENDALE GA: Does estrogen prevent skin aging? Results from the first National Health and Nutrition Examination Survey (NHANES I). *Arch Dermatol* 133:339–342, 1997
  56. SHERWIN BB: Hormones, mood, and cognitive function in postmenopausal women. *Obstet Gynecol* 87(Suppl):20S–26S, 1996
  57. HONJO J, TANAKA K, KASHIWAGI T, URABE M, OKADA H, HAYASHI M, HAYASHI K: Senile dementia-Alzheimer's type and estrogen. *Horm Metab Res* 27:204–207, 1995
  58. KAWAS C, RESNIK S, MORRISON A: A prospective study of estrogen replacement therapy and the risk of developing Alzheimer's disease: The Baltimore Longitudinal Study of Aging. *Neurology* 48:1517–1521, 1997
  59. BARRETT-CONNOR, KRITZ-SILVERSTEIN D: Estrogen replacement therapy and cognitive function in older women. *JAMA* 269:2637–2641, 1993
  60. LIVIO M, MANNUCCHI PM, BIGNANO G: Conjugated estrogens for the management of bleeding associated with renal failure. *N Engl J Med* 315:731, 1986
  61. LOHR JW, SCHWAB SJ: Minimizing hemorrhagic complications in dialysis patients. (abstract) *J Am Soc Nephrol* 2:961, 1991
  62. COLDITZ GA, HANKINSON SE, HUNTER DJ: The use of estrogens and progestins and the risk of breast cancer in postmenopausal women. *N Engl J Med* 332:1589–1593, 1995
  63. BERGKVIST L, ADAMI H-O, PERSSON I: The risk of breast cancer after estrogen and estrogen-progestin replacement. *N Engl J Med* 321:293–297, 1989
  64. JICK H, DERBY LE, MYERS MW, VASILAKIS C, NEWTON KM: Risk of hospital admission for idiopathic venous thromboembolism among users of postmenopausal oestrogens. *Lancet* 348:981–983, 1996
  65. BOSTON COLLABORATIVE DRUG SURVEILLANCE PROGRAM: Surgically confirmed gallbladder disease, venous thromboembolism, and breast tumors in relation to postmenopausal estrogen therapy. *N Engl J Med* 290:15–19, 1974
  66. KESSLER CM, SZYMANSKI LM, SHAMSIPOUR Z, MUESING RA, MILLER VT, LAROSA JC: Estrogen replacement therapy and coagulation: Relationship to lipid and lipoprotein changes. *Obstet Gynecol* 89:326–331, 1997
  67. SCARABIN PY, ALHENC-GELAS M, PLU-BUREAU G, TAISNE P, AGHER R, AIACH M: Effects of oral and transdermal estrogen/progesterone regimens on blood coagulation and fibrinolysis in postmenopausal women. A randomized controlled trial. *Arterioscler Throm Vasc Biol* 17:3071–3078, 1997